

RELICS OF SUBLUMINOUS SUPERNOVAE IN METAL-POOR STARS

TAKUJI TSUJIMOTO¹ AND TOSHIKAZU SHIGEYAMA²
Draft version February 5, 2008

ABSTRACT

The unique elemental abundance pattern of the carbon-rich stars CS29498-043 and CS22949-037 is characterized by a large excess of magnesium and silicon in comparison with iron. This excess is investigated in the context of a supernova-induced star formation scenario, and it is concluded that these stars were born from the matter swept up by supernova remnants containing little iron and that such supernovae are similar to the least-luminous SNe ever observed, SNe 1997D and 1999br. Comparison of the observed abundance pattern in iron-group elements of subluminoous supernovae with those of other supernovae leads to an intriguing implication for explosion, nucleosynthesis, and mixing in supernovae. The observed invariance of these ratios can not be accounted for by a spherically symmetric supernova model.

Subject headings: nuclear reactions, nucleosynthesis, abundances – stars: abundances – stars: individual (CS 29498-043, CS22949-037) – supernovae: general – supernovae: individual (SN 1997D, SN 1999br)

1. INTRODUCTION

Recently, Aoki et al. (2002) revealed a unique elemental abundance pattern in a carbon-rich extremely metal-poor star CS29498-043. The chemical composition of CS29498-043 has a large excess of α -elements such as Mg and Si as well as C and N compared to Fe-group elements. A similar characteristic was previously identified in another C-rich star CS22949-037 (McWilliam et al. 1995; Norris, Ryan, & Beers 2001; Depagne et al. 2002). In CS22949-037, oxygen is also found to be significantly overabundant (Depagne et al. 2002). Since a mixing process at the asymptotic giant branch of stellar evolution enhances exclusively C and N, the lack of Fe is likely to give rise to the enhancements of C, N, Mg and Si (Norris et al. 2001; Aoki et al. 2002). The lack of Fe, of course, must be initially imprinted in stars at formation because the low-mass stars considered here are unable to destroy Fe nuclei in their envelope.

Recent studies on the chemical compositions of metal-poor halo stars have claimed that these stars might have inherited the abundance pattern for the ejecta of the preceding few supernovae (SNe) (Audouze & Silk 1995) or a single SN (Shigeyama & Tsujimoto 1998, ST98 in the following). If this is the case, these stars would have formed from the interstellar matter (ISM) comprising the ejecta of a single SN. Depagne et al. (2002) compared the relative elemental abundances observed in CS22949-037 with those of the theoretical SN models proposed by Woosley & Weaver (1995) and obtained fairly good agreement with a particular model. Furthermore, since the mass of the ISM eventually swept up by the preceding SN is theoretically derived, the observed abundance ratios with respect to H on the surfaces of stars would allow the mass of each element contained in the SN to be estimated (ST98; Tsujimoto & Shigeyama 1998). A comparison of the derived mass of each element with that of theoretical SN models also leads to the conclusion that the apparent enhancement of α -elements in the above two stars is a result of

the deficiency of Fe in the ejecta of the preceding SN (see §2 for quantitative discussion). This indicates that there existed subluminoous SNe with a small amount of ^{56}Ni as a heat source, which decays to ^{56}Fe through ^{56}Co with a half-life of ~ 77 d.

The least luminous SNe ever observed, SN 1997D (Turatto et al. 1998) and SN 1999br (Hamuy 2002, and references therein), are in fact Fe-poor. From analyses of the light curve of SN 1997D (Turatto et al. 1998; Benetti et al. 2001), it has been found that this SN ejects only $\sim 0.002 M_{\odot}$ of ^{56}Ni . The explosion energy E is estimated to be $E \sim 4 \times 10^{50}$ erg and the progenitor mass to be $\sim 25 M_{\odot}$. Hamuy (2002) reported that SN 1999br ejects a similar amount of ^{56}Ni ($\sim 0.0016 M_{\odot}$) with an explosion energy of $\sim 6 \times 10^{50}$ erg and that the ejected mass, which gives the lower bound of the progenitor mass, has been estimated to be $14 M_{\odot}$ from the light curve analyses. It is well known that there is a distinct class of SNe with similar progenitor mass that yield Fe comparable to SN 1987A or slightly higher ($0.07 - 0.15 M_{\odot}$, e.g., ST98). Such SNe will be referred to as *normal* SNe in the following. As mentioned in Turatto et al. (1998), the low Fe mass of SN 1997D may be due to fallback of material onto the collapsed remnant. In this case, Fe as well as other Fe-group elements such as Cr, Mn, Co and Ni produced in the deep interior of the core would fall back. In contrast, lighter elements synthesized in the outer layers of the star would be ejected and be excluded from any fallback. As a result, lighter elements are significantly enhanced with respect to Fe-group elements. The observed elemental abundance patterns of CS29498-043 and CS22949-037 exhibit characteristics that fit this explanation.

In the next section, the features of SNe that induced the formation of the stars CS29498-043 and CS22949-037 will be deduced from the observed elemental abundance pattern in light of the SN-induced star formation scenario proposed by ST98. Subsequently in §3, nucleosynthesis for

¹ National Astronomical Observatory, Mitaka-shi, Tokyo 181-8588, Japan; taku.tsujimoto@nao.ac.jp

² Research Center for the Early Universe, Graduate School of Science, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan; shigeyama@resceu.s.u-tokyo.ac.jp




FIG. 1.— SN II progenitor mass M_{ms} plotted as a function of $[\text{Mg}/\text{H}]$ inside the shell swept up by the SNR. Two cases for SN explosion energy $E = 4 \times 10^{50}$ and 1×10^{51} erg are shown. The observed $[\text{Mg}/\text{H}]$ ratios for stars CS29498-043 (Aoki et al. 2002) and CS22949-037 (Depagne et al. 2002) are indicated by *filled circles* on the curve for $E = 4 \times 10^{50}$ erg. Dashed lines denote the SN progenitor masses corresponding to the preceding SNe for these two stars.

Fe-group elements is discussed based on the implications of the $[\text{Cr}, \text{Mn}, \text{Co}/\text{Fe}]$ ratios observed in these two stars, which is surprisingly at variance with the present theoretical expectations from spherically symmetric SN models.

2. STARS WITH RELICS OF SUBLUMINOUS SUPERNOVAE

The subluminous SNe discovered so far are less energetic varieties of SNe, although theoretically some energetic SNe with high masses such as $35 M_{\odot}$ could also be subluminous (Woosley & Weaver 1995). Such low energy reduces the mass M_{SW} of ISM (mostly hydrogen) eventually swept up by a subluminous SN, obeying the relation $M_{\text{SW}} \propto E^{0.97}$ (ST98). Since the metallicity $[\text{X}/\text{H}]$ of extremely metal-poor stars born from the ISM swept up by a SN remnant (SNR) is inversely proportional to M_{SW} , stars that inherit the ejecta of a subluminous SN should have larger $[\text{X}/\text{H}]$ as long as the other conditions are the same.

As discussed in ST98, the abundance ratios $[\text{Mg}/\text{H}]$ in metal-poor stars can be used as an indicator of the progenitor mass of the SN inducing the formation of the stars, since theoretical models for massive stars predict that the mass of Mg inside a star increases with increasing stellar mass (e.g., Woosley & Weaver 1995; Umeda & Nomoto 2002). In addition, the amount of ejected lighter elements such as Mg can be predicted more precisely than that of heavier elements such as Fe because the ejection of the outer layer is less affected by theoretical uncertainty concerning the SN explosion mechanism (see, however, model Z20A of Woosley & Weaver 1995, for example).

The relationships between the metallicity $[\text{Mg}/\text{H}]$ of stars and the mass M_{ms} of the SN progenitor thus obtained are shown in Figure 1 for $E = 4 \times 10^{50}$ erg (*solid line*) and $E = 1 \times 10^{51}$ erg (*dotted line*), assuming the metallicity $[\text{Mg}/\text{H}]$ of stars born from the shell formed by an SNR is approximated well by the average $[\text{Mg}/\text{H}]$ inside the ISM swept up by the SNR. If CS29498-043 inherited the ejecta of a subluminous SN with explosion energy equal to that of SN 1997D, the SN progenitor mass becomes $25 M_{\odot}$, which intriguingly agrees with that inferred from the light curve analyses of SN 1997D. The SN progenitor

mass for CS22949-037 estimated in the same manner is $M_{\text{ms}} \approx 20 M_{\odot}$. The masses of ejected Fe are estimated to be $0.006 M_{\odot}$ (CS29498-043) and $0.003 M_{\odot}$ (CS22949-037) using the observed $[\text{Fe}/\text{H}]$ ratios for the two stars and assuming $E = 4 \times 10^{50}$ erg. Both of the estimated Fe masses indicate subluminous SNe such as SNe 1997D and 1999br.

Figure 2 shows the enhancement factors of each element defined by the abundance ratios of the element with respect to Fe in the metal-poor stars discussed in this letter relative to those ratios in the ejecta of normal SNe with the corresponding progenitor masses. The abundances of normal SNe have been inferred from the elemental abundance pattern observed for metal-poor stars in the manner described in Tsujimoto & Shigeyama (1998). Thus derived abundance ratios correspond to the average values of metal-poor stars observed by McWilliam et al. (1995) at the metallicity of $[\text{Mg}/\text{H}] \approx -2.3$ and -2.8 , respectively (see Fig. 1). That is, an SN with $E = 1 \times 10^{51}$ erg and $M_{\text{ms}} = 25 M_{\odot}$ will imprint a metallicity $[\text{Mg}/\text{H}] = -2.3$ on the descendant stars, while an SN with the same energy but with $M_{\text{ms}} = 20 M_{\odot}$ corresponds to $[\text{Mg}/\text{H}] = -2.8$. As seen in Figure 2, the abundance of elements heavier than Si in these stars is at least a factor of 5 smaller than the average value of metal-poor stars, which is attributed to fallback onto the remnant in subluminous SNe. A moderate enhancement factor for Si in comparison with Fe-group elements might imply that fallback occurred inside the Si layer. The enhancement factors for Ca through Co have similar values in CS22949-037, while Cr through Fe are similar in CS29498-043. A good coincidence in the enhancement factors is seen especially in CS22949-037, probably due to more accurate abundance determinations. This result suggests that there is little difference in the ratios of Ca, Cr, Mn, or Co to Fe between subluminous SNe and normal SNe with the same progenitor mass, with intriguing implications for nucleosynthesis, explosion, and mixing in SN models.

3. IMPLICATIONS FOR EXPLOSIVE NUCLEOSYNTHESIS AND SUPERNOVA EXPLOSION




FIG. 2.— Abundance ratios of elements with respect to Mg in CS29498-043 (filled circles: Aoki et al. 2002) and CS22949-037 (open circles: Depagne et al. 2002) relative to those in the ejecta of normal SNe with the masses of $25 M_{\odot}$ and $20 M_{\odot}$, respectively. See text for details.

Figure 3 shows the observed correlations between $[\text{Cr}/\text{Fe}]$, $[\text{Mn}/\text{Fe}]$, $[\text{Co}/\text{Fe}]$ and $[\text{Mg}/\text{H}]$ for metal-poor halo stars (McWilliam et al. 1995), CS29498-043, and CS22949-037. As shown in these figures, the observed $[\text{Cr}, \text{Mn}, \text{Co}/\text{Fe}]$ trends are quite clear; the $[\text{Cr}, \text{Mn}/\text{Fe}]$ ratios increase with increasing $[\text{Mg}/\text{H}]$, whereas the $[\text{Co}/\text{Fe}]$ ratio decreases. These clear correlations suggest that the $[\text{Cr}/\text{Fe}]$, $[\text{Mn}/\text{Fe}]$, $[\text{Co}/\text{Fe}]$ ratios are closely correlated with the SN progenitor mass (Tsujiimoto & Shigeyama 1998; Nakamura et al. 1999), while the abundance ratios of α -elements such as Mg, Si, and Ca to Fe do not have clear correlations. It is also clear from the same figure that the observed $[\text{Cr}/\text{Fe}]$, $[\text{Mn}/\text{Fe}]$, $[\text{Co}/\text{Fe}]$ ratios of CS29498-043 and CS22949-037 do not follow the trend presented by the other stars. Five horizontal arrows in this figure indicate how these ratios would shift when these stars inherited the ejecta of an SN with $E = 1 \times 10^{51}$ erg instead of $E = 4 \times 10^{50}$ erg. The lengths of the horizontal shifts correspond to differences in the swept-up hydrogen mass between SNe with $E = 4 \times 10^{50}$ and 1×10^{51} erg. All the arrows pointing to the lines of the best fit for the observational data by χ^2 test, suggesting that the $[\text{Mn}/\text{Fe}]$, $[\text{Cr}/\text{Fe}]$, and $[\text{Co}/\text{Fe}]$ ratios of these two stars do not deviate from those of normal SNe with the same progenitor mass, although for CS29498-043 an uncertainty remains due to the large error in the observed $[\text{Mn}/\text{Fe}]$ ratio and a lack of Co data. As a consequence, there is little difference in the ratios of Cr, Mn, or Co to Fe between subluminal SNe and normal SNe given the same progenitor mass. This turns out to be incompatible with spherically symmetric SNe, as follows.

The elements Mn and Cr are the products of incomplete Si burning, whereas Co is produced by complete Si burning. Fe is produced in both burning processes. The region of complete Si burning is located at the bottom of the ejecta. If an SN explosion were spherically symmetric, different amounts of Fe could only be attributable to different combinations of the mass cut separating the ejecta from the remnant and the explosion energy. Since the mass cut determines the ejected fraction of the region in which complete Si burning takes place, smaller Co/Fe ratios would be expected from SNe with smaller amounts of Fe, corresponding to subluminal SNe. Larger Mn/Fe and Cr/Fe ratios would be expected from these SNe, as shown by

Umeda & Nomoto (2002). Lower explosion energy would suppress complete Si burning and thus reduce both the amount of Fe and the Co/Fe ratio. Therefore, any combination of explosion energy and mass cut in spherically symmetric SNe is incompatible with the observed invariance of the abundance ratios of Fe-group elements to Fe. Since Rayleigh-Taylor instabilities take place after the nuclear reactions freeze out as was shown by a multi-dimensional calculation (Kifonidis et al. 2000), mixing does not alter the relative abundances of Fe-group elements determined in the way mentioned above. The same calculation has also shown that the mixing process is effective. Thus it is expected that fallback that may occur afterwards would hardly change these relative abundances, either. Mixing is unlikely to explain the observed invariance of the relative abundances among SNe with different explosion energies.

We will discuss a possibility that the observed ratios among Fe-group elements for the two stars reflect those in the ISM swept up by an SNR rather than those in individual SN ejecta. Indeed, the gas swept up by the SNR from which stars are formed contains not only heavy elements ejected from the SN itself but also those already present in the ISM. In general, for metal-poor regime such as $[\text{Mg}/\text{H}] \lesssim -2$, the stellar abundance pattern is essentially determined by the SN yield due to a negligible contribution from the ISM (ST98; Tsujiimoto, Shigeyama, & Yoshii 1999). A lack of iron-group elements in the ejecta of a sub-luminous SN could enhance contribution from the ISM to the stellar abundance of iron-group elements. Since the $[\text{Cr}/\text{Fe}]$, $[\text{Mn}/\text{Fe}]$, $[\text{Co}/\text{Fe}]$ ratios in the ISM should be equal to nearly zero, which corresponds to the plateau value seen in halo stars with $-2 < [\text{Fe}/\text{H}] < -1$, the two stars discussed in this letter should show these abundance ratios nearly equal to the solar values if this is the case. However, the observed ratios shown in Figure 3 do not support this possibility. Therefore it is likely that these stars contain little heavy elements from the ISM. Furthermore, combinations of these ratios predicted from spherically symmetric SN models with those in the ISM can not make the observed ones.

Stellar rotation might be a factor to produce a variety in energies of supernova explosions with the same progenitor mass. It is sometimes suggested that a rapid rotation leads to a jet-like explosion (e.g., Woosley 1993). In fact, it has




FIG. 3.— (a) The correlation between $[\text{Cr}/\text{Fe}]$ and $[\text{Mg}/\text{H}]$ for metal-poor stars observed by McWilliam et al. (1995), CS29498-043 (Aoki et al. 2002) and CS22949-037 (Depagne et al. 2002). The solid line shows the χ^2 fit to the observed points. Arrows indicate how the $[\text{Mg}/\text{H}]$ ratios for these stars would shift if these stars were born from the SNRs originating from SNe with $E = 1 \times 10^{51}$ ergs. (b) The same as (a) but for Mn. (c) The same as (a) but for Co.

been a preferred explanation for matter mixing suggested by the emission line profiles observed for SN 1987A (Nagataki 2000) and SN 1998bw (Maeda et al. 2002), both of which are core-collapse SNe. Furthermore, the optical spectropolarimetry of several core-collapse SNe suggests highly asymmetric explosion mechanisms such as the formation of bipolar jets (Wang et al. 2001). Detailed numerical studies on nucleosynthesis and ejection mechanism in jet-like explosions might be required to explain the observed trend of Fe-group elements in halo field stars.

An alternative solution may be in a hot bubble formed by neutrino emission. If neutrinos from the nascent neutron star deposit substantial energy to make a hot bubble behind the blast wave, the hot bubble may carry exclusively the product of complete Si burning to the outer layer by buoyancy. A numerical simulation by Wilson & Mayle (1988) has shown that this mechanism leads to a less energetic SN. This may suppress fallback of Co and Fe in subluminal SNe, though other numerical studies have shown that a hot bubble does not result in a SN explosion (e.g., Bruenn, Mezzacappa, & Dineva 1995). The current understanding of the involved physics is, however, still too premature to extract a definite conclusion on this issue.

4. CONCLUSIONS

The recently discovered new class of carbon-rich stars including CS29498-043 and CS22949-037, which exhibit a significant enhancement of α -elements with respect to Fe-group elements, has been shown in this study to inherit the elemental abundance pattern of a precursor SNe that ejected a small but adequate amount of Fe with low explosion energy. These SNe are found to resemble the two recently discovered subluminal SNe 1997D and 1999br in terms of the ejected Fe mass, explosion energy, and progenitor mass. Subluminal supernovae may therefore be studied more easily by looking at the surface of nearby metal-poor stars than by searching for similar occurrences elsewhere.

The present study revealed that there is little difference in the $[\text{Cr}/\text{Fe}]$, $[\text{Mn}/\text{Fe}]$, $[\text{Co}/\text{Fe}]$ ratios between subluminal SNe and normal SNe given the same progenitor mass, which has intriguing implications in nucleosynthesis and explosion in SN models. The observed facts are apparently at odds with the prediction by spherically symmetric SN explosion models.

We are grateful to the anonymous referee for making useful comments. This work was supported in part by a

Grant-in-Aid for Scientific Research (11640229, 12640242) and Technology of Japan.
from the Ministry of Education, Culture, Sports, Science,

REFERENCES

- Aoki, W., Norris, J. E., Ryan, S. G., Beers, T. C., & Ando, H. 2002, *ApJ*, 576, L141
 Audouze, J., & Silk, J. 1995, *ApJ*, 451, L49
 Benetti, S., Turatto, M., Balberg, S., Zampieri, L., Shapiro, S. L., Cappellaro, E., Nomoto, K., Nakamura, T., Mazzali, P. A., Patat, F. 2001, *MNRAS*, 322, 361
 Bruenn, S. W., Mezzacappa, A., & Dineva, T. 1995, *Phys. Rep.* 256, 69
 Depagne, E. et al. 2002, *A&A*, 390, 198
 Hamuy, M. 2003, *ApJ*, preprint doi:10.1086/34689
 Kifonidis, K., Plewa, T., Janka, H.-Th., & Müller, E. 2000, *ApJ*, 531, L123
 Maeda, K., Nakamura, T., Nomoto, K., Mazzali, P. A., Patat, F., & Hachisu, I. 2002, *ApJ*, 565, 405
 McWilliam, A., Preston, G. W., Sneden, C., & Searle, L. 1995, *AJ*, 109, 2757
 Nagataki, S. 2000, *ApJS*, 127, 141
 Nakamura, T., Umeda, H., Nomoto, K., Thielemann, F.-K., & Burrows, A. 1999, *ApJ*, 517, 193
 Norris, J. E., Ryan, S. G., & Beers, T. C. 2001, *ApJ*, 561, 1034
 Shigeyama, T., & Tsujimoto, T. 1998, *ApJ*, 507, L135 (ST98)
 Tsujimoto, T., & Shigeyama, T. 1998, *ApJ*, 508, L151
 Tsujimoto, T., Shigeyama, T., & Yoshii, Y. 1999, *ApJ*, 519, L63
 Turatto, M. et al. 1998, *ApJ*, 498, L129
 Umeda, H., & Nomoto, K. 2002, *ApJ*, 565, 385
 Wang, L., Howell, D. A., Höflich, P., & Wheeler, J. C. 2001, *ApJ*, 550, 1030
 Wilson, J. R., & Mayle, R. W. 1988, *Phys. Rep.* 163, 63
 Woosley, S. E. 1993, *ApJ*, 405, 273
 Woosley, S. E., & Weaver, T. A. 1995, *ApJS*, 101, 181